


## 1 Markov Chain Terminology

In this question, we will walk you through terms related to Markov chains.

Note 22

1. (Irreducibility) A Markov chain is irreducible if, starting from any state  $i$ , the chain can transition to any other state  $j$ , possibly in multiple steps.
2. (Periodicity)  $d(i) := \gcd\{n > 0 \mid P^n(i, i) = \mathbb{P}[X_n = i \mid X_0 = i] > 0\}$ ,  $i \in \mathcal{X}$ . If  $d(i) = 1 \forall i \in \mathcal{X}$ , then the Markov chain is aperiodic; otherwise it is periodic.
3. (Matrix Representation) Define the transition probability matrix  $P$  by filling entry  $(i, j)$  with probability  $P(i, j)$ .
4. (Invariance) A distribution  $\pi$  is invariant for the transition probability matrix  $P$  if it satisfies the following balance equations:  $\pi = \pi P$ .



(a) For what values of  $a$  and  $b$  is the above Markov chain irreducible? Reducible?

(b) For  $a = 1$ ,  $b = 1$ , prove that the above Markov chain is periodic.

(c) For  $0 < a < 1$ ,  $0 < b < 1$ , prove that the above Markov chain is aperiodic.

(d) Construct a transition probability matrix using the above Markov chain.

(e) Write down the balance equations for this Markov chain and solve them. Assume that the Markov chain is irreducible.

## 2 Can it be a Markov Chain?

Note 22

- (a) A fly flies in a straight line in unit-length increments. Each second it moves to the left with probability 0.3, right with probability 0.3, and stays put with probability 0.4. There are two spiders at positions 1 and  $m$  and if the fly lands in either of those positions it is captured. Given that the fly starts between positions 1 and  $m$ , model this process as a Markov Chain.

- (b) Take the same scenario as in the previous part with  $m = 4$ . Let  $Y_n = 0$  if at time  $n$  the fly is in position 1 or 2 and let  $Y_n = 1$  if at time  $n$  the fly is in position 3 or 4. Is the process  $Y_n$  a Markov chain?

### 3 Allen's Umbrella Setup

Note 22

Every morning, Allen walks from his home to Soda, and every evening, Allen walks from Soda to his home. Suppose that Allen has two umbrellas in his possession, but he sometimes leaves his umbrellas behind. Specifically, before leaving from his home or Soda, he checks the weather. If it is raining outside, he will bring exactly one umbrella (that is, if there is an umbrella where he currently is). If it is not raining outside, he will forget to bring his umbrella. Assume that the probability of rain is  $p$ .

- (a) Model this as a Markov chain. What is  $\mathcal{X}$ ? Write down the transition matrix.
- (b) What is the transition matrix after 2 trips?  $n$  trips? Determine if the distribution of  $X_n$  converges to the invariant distribution, and compute the invariant distribution. Determine the long-term fraction of time that Allen will walk through rain with no umbrella.